

# SC



**UNEP**/POPS/POPRC.6/13/Add.2

---

Distr.: General  
3 December 2010

Arabic  
Original: English



/ -

.UNEP/POPS/POPRC.6/10

/

.....		-
.....		-
.....		-
.....		-
.....		-
.....		-
.....		-
.....		-
.....	- -	
.....	- -	
.....	- -	
.....		-
.....	- -	
.....	- -	
.....	- -	
.....		-
.....	- -	
.....	- -	
.....		-
.....	- -	
.....	- -	
.....	- -	
.....	- -	
.....	- -	
.....	- -	
.....	- -	
.....		-
.....		-
.....		
.....		

)

( )

(





/

. /





:

		
		
- :	- - : - - :	- - : - - :

:

-	:
- - - -	:
- - - -	:
-	:
C <sub>12</sub> H <sub>18</sub> Br <sub>6</sub>	:
,	:
Cyclododecane, hexabromo; HBCD; Bromkal 73-6CD; Nikkafainon CG 1; Pyroguard F 800; Pyroguard SR 103; Pyroguard SR 103A; Pyrovatex 3887; Great Lakes CD-75P™; Great Lakes CD-75; Great Lakes CD75XF; Great Lakes CD75PC (compact); Dead Sea Bromine Group Ground FR 1206 I-LM; Dead Sea Bromine Group Standard FR 1206 I-LM; Dead Sea Bromine Group Compactified FR 1206 I-CM.	/

<p>( )</p>	<p>( )</p>
------------	------------

( ) :

	$C_{12}H_{18}Br_6$	
	,	
( )	- -	
	. - : - : - :	
( )	)	
( )	(	
( )	/ ,	
( )	/ ,	
( )	( ) - × ,	
	( )	
( )	, =	
( )	( )	/
	( ) , ± ,	
	( ) , ± ,	
	( ) , ± ,	
	/ × ,	
	)	
	( /	
	-	
	/ , =	
	, = /	



:

.( )

( )

	*( / )		
( )	, ± ,		-
	, ± ,		-
	, ± ,		-
	,		( )
( )	,		-
	,		-
	,		-
	,		( )
( )	** , ± ,		-

\*\* \*

-

-

(UNEP/POPS/POPRC.5/10 ) /

.( / - )

-

-

“ ”  
“ ”

)

-

.(

-

(ECE/EB.AIR/WG.5/2009/7 )

/

/

.( )

( )

)

.(

-

)

.(

-

( )

/ .( )

)

)( / ( )

:

.(

-

.( )

-

-

/

-

-

-

--

-

)

.(

-

.( )

( )

.( )

.( )

/

( )

.( )  
 ) ( - ) : ( ) ( ) ( - -  
 .( )  
 )  
 -  
 .( ) .  
 )  
 .( )  
 -  
 )  
 )  
 .( )  
 )  
 ) - .( )  
 ) .( )  
 ) .( )  
 ) .( )  
 )  
 ) ( ) .( )  
 .  
 ) ( ) - )  
 )  
 )  
 )  
 ) ( )  
 .  
 ) ( )  
 )

.( )  
( )  
) ( ) .( )  
( )

.( -

( )

( )

)  
(

: / : / : )  
 ( / : / : ) ( / )  
 )  
 (

./ /  
 /  
 /  
 /  
 /  
 /

./ ( )  
 )  
 ( / ) ( / )  
 ) ( )  
 ( / , )  
 .(

./ ( )  
 ( )  
 )  
 / :  
 .( /



(

. ,

( )

.( )

-

.( )

.

.( )

.( )

-

)

.( )

.( ) (

-

--

-

( - )

.

.( )

( )



/ , )

.( /

( )

( )

.( )

( )

( )

( )

( )

.( )

( / )

( / )

)

( )

.(

.( )

(-/+)

(-/+)

(-/+)

( ) ( )

)

)  
( )

( )

( / )

)

.(

( )  
(*Pimephales promeas*) ( )

(*Oncorhynchus mykiss*)

/ , ±

( ) / , , .( )

( ) ( )

.( )

( )

- / -

- / , /

.( ) / ,

( ) -

( ' ' ' )

( ) -

<< <

)  
 / )  
 ( ) ( /  
 .  
 ( - : - : - : )  
 .( , : , : , : )  
 ( )  
 ( ) <sup>14</sup>C

.( )

- :

( , > ) ,  
 .( )

.( )

( )

( / , )

(*Boreogadus saida*)

.(*Gammarus wilkitzkii*)

( , )

.( , )

/ / ( )

(*Pusa hispida*)

. ( ) (*Ursus maritimus*)

( ) -

(*Phoca vitulina*)

(*Phocoena phocoena*)

. (*Anguilla Anguilla*)

( : ) :

( : : ) ::

( )

)

(

(*Carassius carassius*)

(*Littorina littorea*)

( ) -

( , > ) ,

) ,

)

. (

(*Sebastes mentella*)

(*Pandalus borealis, Hymenodora glacialis*)

*Delphinapterus* )

(*Monodon monoceros*)

(*Boreogadus saida*)

(*leucas*)

. (*Odobenus rosmarus*)

(*Mya truncata, Serripes groenlandica*)

( )

)

( ) .(

) (

- -

-

-

.( - ) ( , )

×

.( )

.( )

( ) -

× , - / × , - / × , /

( )

× , - / × ( , ± ) ,

- /

.)

.(

( ) -

)

-

. ( - )  
)  
(  
.  
-  
.  
-  
,  
( )  
( )  
.  
.  
-  
"  
"  
/  
( )  
/  
( )  
( )  
( , - , )  
"  
"  
-  
"  
"  
)  
(

“ ”

.

-

( ) .( )

.

.( )

(“ ” / , “ ” / , )

, )

(“ ” / , “ ” /  
.( )

.( )

.( )

( / - )

-

( ) / -

/ , )

-

.(

.( ) -

-

--

-

( )

( )

.

.

*(Delphinus delphis)*

.

)

( /

/



.( / )  
 /  
 /  
 .( ) /  
 /  
 /  
 .( )  
 - ) (*Accipiter nisus*) ( / - ) (*Falco peregrinus*)  
 ( / )  
 ( )  
 ( *Sterna hirundo* )  
 / -  
 .( )  
 -  
 .( )  
 )  
 .( ) / , ( /  
 / / /  
 - )  
 ( ) .( ) ( /  
 / , . ( - ) (*pelamis Katsuwonus*)  
 /  
 ( )  
 (*Ctenopharyngodon idella*) /  
 .(*Siniperca chuatsi*) /

)  
 .( )  
 .( )  
 ( )  
 (*Delphinapterus leucas*)  
 / - ,  
 (*Ursus maritimus*) ( )  
 / )  
 .( )  
 ( )  
 (*Pagophila eburnea*)  
 .( )  
 .  
 .(www.iucnredlist.org/)  
 ( ) /  
 (*Larus marinus*) (*Larus hyperboreus*)  
 -  
 .  
 . (*Larus hyperboreus*)  
 / , (*Larus hyperboreus*)  
 / , /  
 / , , / , ,  
 / - - (*Phalacrocorax carbo*)  
 - - .( )  
 (*Larus hyperboreus*)  
 ( )



( - ) ( ) -

.( )  
( ) -

*(Zalophus californianus)*

( )  
.( - )

( / - )  
/  
/

)  
.( /

.(

)  
.( )

(  
.( )

.  
.  
( ) -  
.  
( )  
/  
/  
) /  
( ) ( ) /  
( ) ( ) ( ) -  
( ) ( ) ( ) :  
( ) ( )  
)  
( ) .(  
/  
/  
/  
( ) / , ,  
( / , )  
( ) ( )  
/



/ .(

.

.( )

)

.

.(

( )

( )

.

.( )

( )

:

.

.( )

---

-

:

)

.( )

.( )

( )

.

/ - )

, , )

/

.(

( )

/

.( )

( )

-

( )

.

'

/

/

/

( / ) ( )  
 ( / ) ( / )  
 .( / )  
 .( )  
 / , .( )  
 )  
 ) ( / , .( )  
 ) )  
 ) ) .( )  
 ) / , .( )  
 .( / ) ( / ) ( )  
 .( / ) ( / ) ( )  
 ( )  
 )  
 .( ) .( )  
 ( )  
 .( )  
 ( )  
 )  
 .( )





( )

.( ) / /

( )

)

.(

. / / /

.( ) /

-

/ / /

) , / / /

.(

/ /

.( )

.( ) " " / / ,

-

( )

( )

.(

-

)

-

.(

.( )  
-  
.  
.( )  
--  
-  
)  
.( )  
.  
.( )  
,  
Skeletonema ) ( Daphnia magna ) /  
.( / ) (costatum  
.( / )  
(Lumbriculus variegates)  
.( )  
-  
.  
.( )  
/ , ± , , ± , , ± ,  
.( ± )  
(P450 CYP1A )  
." "

.( ) / ( )  
 .  
 .  
 .( )

*(Platichthys flesus)*

.( )  
 / ) ( / )  
 + + , + , + , ( ) + : ( + +  
 .  
 ) (T3)  
 . ( -  
 ( )

*(Danio rerio)*

' ' ' ' ' ' /  
 ( / ) ( / )  
 )  
 ( . / ,

( - - - )  
 ( - - ) .  
 .  
 - )  
 ( ) .(  
 ( / , , )  
 ( / , ≥ )  
 ( / , , )  
 .( / , ≤ ) (Hsp70)  
 (*Gobiocyprinus rarus*) ( )  
 )  
 ( ) ( )  
 .( ) /  
 / ( ) )  
 .  
 .( )  
 .( ) (*Macoma balthica*)  
 - - -  
 ( )  
 ( ) (*Eisenia fetida*)  
 , /  
 /  
 /  
 .( )  
 < ( )  
 (*Allium cepa*) (*Cucumis sativa*) (*Zea mays*) ( )  
 /

*(Lycopersicon esculentum)*

*(Glycine max)*

*(Lolium perenne)*

/

≤

.( )

--

-

)

.(

) < < << :

/

.(

,

/

,

(

/

)

/

/

)

.(

)

(

) (*Falco sparverius*)

/

.(

.

/

.(

)

.(

)

/

)

( )

(

/

-

:

( )

/

-

:

( )

/

-

:

( )

(

)

( ) (

)

/

-

:

.(

)

/

,

:

-

*(Coturnix coturnix japonica)*

:

)

(

( / / , )

/ , )

.( ) ( /

-(

.( ) /

(UGT1A9) (CYP3A37) (CYP2H1)

)

( ) .(

CYP2H1) ( )

(UGT1A9) (CYP3A37

(UGT1A1) (UGT1A9)

(T4)

)

( ) .(

(UGT1A9)

/

--

-

)

:

.( )

-

.)

)

- .(

.)

/

)

.( - ) .(

)

(

.

.

)

.(

.)

.)

-(

( )

)

-(

(

)



)  
.(  
/ , / , ) ( )  
.  
( )  
.  
( )  
.( )  
/  
) Ca<sup>2+</sup>  
.(  
( ) -  
.  
.( )  
,  
( ) / ≥  
(T3)  
(T3)  
)  
.(



.( ) ( )  
.  
.  
( ) / /  
.  
--  
-  
( )  
)  
( )  
( )  
( ) ( ) ( ) ( )  
( ) ( ) ( )  
--  
-  
( ) (prey)  
( ) (predators) /  
) /  
(

.( ) / , /

.( )

( ) ( )

/

- :

) / - : / .( /

' / , ( / ) / / , ) ( / , ) .( ) (

) ) - .(

) / , - , (

.(UNEP/POPS/POPRC.6/INF/25

( ) ( )

.( )

( )

( / )

( )

( )

( ) / , - , (Larus hyperboreus)



/

)

.(

:

	( / )	
<p>.</p>		
<p>./ ( ) ( ) ) (</p>		
<p>.</p>		
<p>./ (Daphnia magna) (Skeletonema costatum) : . . - ) / / - ) / / ' ) ( / / ) / / ) ( / / ) ( / / ) .</p>		

	( / )	
/ .( / )	•	

-

-

.( )



:

Abdallah MA, Harrad S. Personal exposure to HBCDs and its degradation products via ingestion of indoor dust. *Environ Int.* 2009;35(6):870-6.

Abdallah MA, Harrad S, Covaci A. Hexabromocyclododecanes and tetrabromobisphenol- A in indoor air and dust in Birmingham, U.K: implications for human exposure. *Environ Sci Technol.* 2008a;42(18):6855-61.

Abdallah Mohamed AE, Harrad S, Ibarra C, Diamond M, Melymuk L, Robson M, Covaci A. Hexabromocyclododecanes in indoor dust from Canada, the United Kingdom, and the United States. *Environ Sci Technol.* 2008b;42(2):459-64

{ACCBFRIP} (American Chemistry Council Brominated Flame Retardant Industry Panel). HPV data summary and test plan for the hexabromocyclododecane (HBCD) CAS no. 3194556. Arlington, VA, USA. Report 2001, 23 pp.

Albemarle Corporation. 1994; Baton Rouge, L.A. USA. As cited in: European Commission, 2008

Albemarle Corporation. 2000. Saytex 9006L Flame Retardant. Baton Rouge. Louisiana: Albemarle Corporation. 2 pp.

Allchin CR, Morris S. Hexabromocyclododecane (HBCD) diastereoisomers and brominated diphenyl ether congener (BDE) residues in edible fish from the rivers Skerne and Tees, U.K. *Organohalogen Compd.* 2003, 61, 41-44.

Antignac JP, Cariou R, Maume D, Marchand P, Monteau F, Zalko D, Berrebi A, Cravedi JP, Andre F, Le Bizec B. Exposure assessment of fetus and newborn to brominated flame retardants in France: preliminary data. *Mol Nutr Food Res.* 2008;52(2):258-65.

{AOPWIN} Atmospheric Oxidation Program for Windows {Estimation Model} Version 1.92. Washington (DC): U.S. Environmental Protection Agency, Office of Pollution Prevention and Toxics; Syracuse (NY): Syracuse Research Corporation. 2000. Available from: [www.epa.gov/oppt/exposure/pubs/episuite.htm](http://www.epa.gov/oppt/exposure/pubs/episuite.htm)

{AMAP} (Arctic Monitoring and Assessment Programme). Arctic Pollution 2009. Report. 83pp.

{ABC} (ABC Laboratories, Inc). Effect of hexabromocyclododecane on the survival and reproduction of the earthworm, *Eisenia fetida*. ABC study No. 47222, ABC Laboratories, Inc. and Wildlife International Ltd, Columbia, Missouri and Easton, Maryland, USA, 2003. Authors: Aufderheide J, Jones A, MacGregor JA and Nixon WB. 94 pp.

Backus S, Batchelor M, Alaee M, Ueno D, and Hewitt LM. Isomer-specific determination of hexabromocyclododecane in abiotic and biotic samples by high performance liquid chromatography/atmospheric pressure photoionization tandem mass spectrometry. *Organohalogen Compd* 2005;240-243.

Bahm, K and Khailil, MAK. (A new model of tropospheric hydroxyl radical concentrations. *Chemosphere* 2004;54: 143-166.

Barres BA, Hart IK, Coles HS, Burne JF, Voyvodic JT, Richardson WD, Raff MC. Cell death and control of cell survival in the oligodendrocyte lineage. *Cell.* 1992;70(1):31-46.

Bergander L., Kierkegaard A., Sellström U, Wideqvist U, and de Wit, C. Are brominated flame retardants present in ambient air? Conference Proceeding, 6th Nordic Symposium of Organic Pollutants. 17-9-1995. [

BFR 2009a, Mattioli L, Fernie KJ, Bird D, Ritchie IJ, Shutt LJ, Letcher RJ. Hexabromocyclododecane (HBCD) isomers in American kestrels (*Falco sparverius*) exposed via the diet to a technical HBCD formulation; Uptake, depuration and bioisomerization. Poster presentation. 11th Annual Workshop on Brominated Flame Retardants (BFR 2009), Ottawa, Canada, 2009.

BFR 2009b, Martenson SC, Bird DM, Letcher R, Shutt L, Fernie K. Behavioural and reproductive changes in American kestrels (*Falco sparverius*) exposed to technical hexabromocyclododecane (HBCD) at environmentally relevant concentrations. Oral presentation. 11th Annual Workshop on Brominated Flame Retardants (BFR 2009), Ottawa, Canada, 2009.

{BIOWIN} Biodegradation Probability Program for Windows {Estimation Model} Version 4.02. Washington (DC): US Environmental Protection Agency, Office of Pollution Prevention and Toxics; Syracuse (NY): Syracuse Research Corporation. 2000. Available from: [www.epa.gov/oppt/exposure/pubs/episuite.htm](http://www.epa.gov/oppt/exposure/pubs/episuite.htm)

- de Boer J, Leslie HA, Leonards PEG, Bersuder P, Morris S, Allchin CR. 2004. Screening and time trend study of decabromodiphenylether and hexabromocyclododecane in birds. Abstract. The Third International Workshop on Brominated Flame Retardants, June 6-9, Toronto, Canada. pp. 125-128.
- Bogdal C, Schmid P, Kohler M, Müller CE, Iozza S, Bucheli TD, Scheringer M, Hungerbühler K. Sediment record and atmospheric deposition of brominated flame retardants and organochlorine compounds in Lake Thun, Switzerland: lessons from the past and evaluation of the present. *Environ Sci Technol*. 2008;42(18):6817-22.
- Boulware MI, Mermelstein PG. The influence of estradiol on nervous system function. *Drug News Perspect*. 2005;18(10):631-7. Polychlorinated dibenzo-p-dioxins, and -furans, dioxin-like polychlorinated biphenyls, brominated flame retardants, perfluorinated alkyl substances, pesticides, and other compounds. *J Environ Monit*. 2007;9(5):465-72. DOI: 10.1039/b617103f
- Brandsma SH, Van der Ven LT, de Boer J, Leonards PE. Identification of hydroxylated metabolites of hexabromocyclododecane in wildlife and 28-days exposed Wistar rats. *Environ Sci Technol*. 2009;43(15):6058-63.
- Brown TN, Wania F. 2008. Screening chemicals for the potential to be persistent organic pollutants: A case study of Arctic contaminants. *Environ Sci Technol* 42(14): 5202-5209.
- {BSEF} (Bromine Science and Environmental Forum). Assessing the long-range transport potential of tetrabromobisphenol A and hexabromocyclododecane using several multimedia transport models. A report to BSEF by WECC, Wania Environmental Chemists Corp. 2003. Author: Wania F. pp 13 pp.
- {BSEF} (Bromine Science and Environmental Forum). About Hexabromocyclododecane (HBCD). 2006. <http://www.bsef.com/our-substances/hbcd/about-hbcd/> (accessed January 2008).
- {BSEF} (Bromine Science and Environmental Forum). About Hexabromocyclododecane (HBCD). 2010. <http://www.bsef.com/our-substances/hbcd/about-hbcd/> (accessed June 2010).
- Chakraborti A, Gulati K, Ray A. Estrogen actions on brain and behavior: recent insights and future challenges. *Rev Neurosci*. 2007;18(5):395-416.
- Chemicals Inspection and Testing Institute, Final Report - Bioconcentration Study of Hexabromocyclododecane in Carp Conducted with 1,2,5,6,9,10-Hexabromocyclododecane, (1995)  
[http://www.meti.go.jp/policy/chemical\\_management/english/files/Bioconcentration\\_study\\_HBCD.pdf](http://www.meti.go.jp/policy/chemical_management/english/files/Bioconcentration_study_HBCD.pdf) (accessed on 11 March 2010).
- Chemicals Inspection and Testing Institute, Final Report - Biodegradation Study of Hexabromocyclododecane Conducted with 1,2,5,6,9,10-Hexabromocyclododecane, (1990)  
[http://www.meti.go.jp/policy/chemical\\_management/english/files/Biodegradation\\_study\\_of\\_HBCD.pdf](http://www.meti.go.jp/policy/chemical_management/english/files/Biodegradation_study_of_HBCD.pdf) (accessed on 11 March 2010).
- Colles A, Koppen G, Hanot V, Nelen V, Dewolf MC, Noël E, Malisch R, Kotz A, Kypke K, Biot P, Vinkx C, Schoeters G. Fourth WHO-coordinated survey of human milk for persistent organic pollutants (POPs): Belgian results. *Chemosphere*. 2008;73(6):907-14.
- Conley A, Hinshelwood M. Mammalian aromatases. *Reproduction* 2001;121:685-695.
- Covaci A, Roosens L, Dirtu AC, Waegeneers N, Van Overmeire I, Neels H, Goeyens L. Brominated flame retardants in Belgian home-produced eggs: levels and contamination sources. *Sci Total Environ*. 2009; 407(15):4387-96.
- Covaci A, Gerecke AC, Law RJ, Voorspoels S, Kohler M, Heeb NV, Leslie H, Allchin CR, de Boer J. Hexabromocyclododecanes (HBCDs) in the environment and humans: a review. *Environ Sci Technol*. 2006; 40(12):3679-88.
- Crump D, Egloff C, Chiu S, Letcher RJ, Chu S, Kennedy SW. Pipping success, isomer-specific accumulation, and hepatic mRNA expression in chicken embryos exposed to HBCD. *Toxicol Sci*. 2010;115(2):492-500
- Crump D, Chiu S, Egloff C, Kennedy SW. Effects of hexabromocyclododecane and polybrominated diphenyl ethers on mRNA expression in chicken (*Gallus domesticus*) hepatocytes. *Toxicol Sci*. 2008;106(2):479-87.
- Davis JW, Gonsior S, Markham DA, Friederich U, Hunziker RW, Ariano J. Biodegradation and product identification of [<sup>14</sup>C]hexabromocyclododecane in wastewater sludge and freshwater aquatic sediment. *Environ Sci Technol* 2006a; 40: 5395-5401.
- Davis JW. The aerobic biodegradability of trans, trans, trans-1,5,9-[<sup>14</sup>C]Cyclododecatriene in a modified ready biodegradation test. 2006b. [No other publication information available].

- Davis JW, Gonsior SJ, Marty G, Ariano JM. The transformation of hexabromocyclododecane in aerobic and anaerobic soils and aquatic sediments. *Water Research* 2005;39: 1075-1084.
- Davies AM. Regulation of neuronal survival and death by extracellular signals during development. *EMBO J.* 2003 2;22(11):2537-45.
- Deng J, Yu L, Liu C, Yu K, Shi X, Yeung LW, Lam PK, Wu RS, Zhou B. Hexabromocyclododecane-induced developmental toxicity and apoptosis in zebrafish embryos. *Aquat Toxicol.* 2009;93(1):29-36.
- Desjardins D, MacGregor JA and Krueger HO. Hexabromocyclododecane (HBCD): A 72-hour toxicity test with the marine diatom (*Skeletonema costatum*), Final report. Wildlife International, Ltd., Easton, Maryland, USA. 2004; 66 pp.
- Dingemans MM, Heusinkveld HJ, de Groot A, Bergman A, van den Berg M, Westerink RH. Hexabromocyclododecane inhibits depolarization-induced increase in intracellular calcium levels and neurotransmitter release in PC12 cells. *Toxicol Sci.* 2009;107(2):490-7.
- Dioxin 2010a, Esslinger S, Becker R, Jung C, Schröter-Kermani C, Nehls I. Time courses of HBCD levels and enantiomeric signatures in Herring gull eggs from the German coast. *Dioxin 2010*, San Antonio, Texas, U.S.A.
- Dioxin 2010b, Fernie KJ, Marteinson SC, Bird DM, Ritchie IJ, Letcher RJ. Changes in reproduction and behavior of American kestrels (*Falco sparverius*) associated with exposure to environmentally relevant concentrations of technical HBCD. *Dioxin 2010*, September 2010, San Antonio, Texas, U.S.A.
- Dioxin 2010c, Marteinson SC, Kimmins S, Letcher RJ, Bird DM, Ritchie IJ, Fernie KJ. Evidence of endocrine disruption and testicular changes in male American kestrels (*Falco sparverius*) exposed to technical HBCD. *Dioxin 2010*, San Antonio, Texas, U.S.A.
- {DOW} (The Dow Chemical Company). Investigation of the biodegradation of <sup>14</sup>C, sediment and soil. Toxicology and Environmental Research and Consulting, The Dow Chemical Company, Midland, Michigan, USA. 2004. Authors: Davis JW, Gonsior SJ, Markham DA, Marty GT. 113 pp.
- Driffield M, Harmer N, Bradley E, Fernandes AR, Rose M, Mortimer D, Dicks P. Determination of brominated flame retardants in food by LC-MS/MS: diastereoisomer-specific hexabromocyclododecane and tetrabromobisphenol A. *Food Addit Contam Part A Chem Anal Control Expo Risk Assess.* 2008;25(7):895-903.
- {EBFRIP} (European Brominated Flame Retardant Industry Panel). A Survey of HBCD Potential Emissions in Europe 2008-2009. Fact sheet, 2009a.
- {EBFRIP} (European Brominated Flame Retardant Industry Panel). An evaluation of hexabromocyclododecane (HBCD) for Persistent Organic Pollutant (POP) properties and the potential for adverse effects in the environment. Report submitted to EBFRIP. Authors: Arnot J, McCarty L, Armitage J, Toose-Reid L, Wania F, Cousins I. 2009, 2009b. 214 pp.
- EBFRIP (European Brominated Flame Retardant Industry Panel). HBCD in polystyrene foams: product safety assessment. PlasticsEurope, EXIBA, EBFRIP. Confidential industry report. 2009c.
- {ECHA} (European Chemicals Agency). Data on manufacture, import, export, uses and releases of HBCDD as well as information on potential alternatives to its use. 2008a, 108 pp
- {ECHA} (European Chemicals Agency). Member state committee support document for identification of hexabromocyclododecane and all major diastereoisomers as a substance of very high concern. 2008b, 43 pp. Available at: [http://echa.europa.eu/chem\\_data/authorisation\\_process/candidate\\_list\\_table\\_en.asp](http://echa.europa.eu/chem_data/authorisation_process/candidate_list_table_en.asp)
- {ECT} Hexabromocyclododecane (HBCD): Effects on soil microorganisms. ECT study number: AU1BB. 2007. Author: Förster B. 20 pp
- Eljarrat E, Guerra P, Martínez E, Farré M, Alvarez JG, López-Teijón M, Barceló D. Hexabromocyclododecane in human breast milk: levels and enantiomeric patterns. *Environ Sci Technol.* 2009 ;43(6):1940-6.
- Ema M, Fujii S, Hirata-Koizumi M, Matsumoto M. Two-generation reproductive toxicity study of the flame retardant hexabromocyclododecane in rats. *Reprod Toxicol.* 2008;25(3):335-51.
- {EMEP} EMEP contribution to the preparatory work for the review of the CLRTAP protocol on persistent organic pollutants. New substances: Model assessment of potential for long-range transboundary atmospheric transport and persistence of Hexabromocyclododecane (HBCDD). EMEP MSC-E Information Note 4/2009. Authors: Vulykh N, Rozovskaya O, Shatalov V.

- {Environment Canada} Data collected pursuant to Section 71 of the Canadian Environmental Protection Act, 1999 and in accordance with the published notice "Notice with Respect to Certain Substances on the Domestic Substances List (DSL), Canada Gazette, Vol. 135 #46". 2001. 21 pp.
- Eriksson P, Fisher C, Wallin M, Jakobsson E and Fredriksson A. Impaired behaviour, learning and memory, in adult mice neonatally exposed to hexabromocyclododecane (HBCDD). *Environ Toxicol Pharmacol* 2006; 21: (3): 317-322.
- {European Commission} Risk assessment hexabromocyclododecane, CAS-No.: 25637-99-4, EINECSNo.: 247-148-4, Final Report May 2008. 492 pp.
- Fery Y, Buschauer I, Salzig C, Lang P, Schrenk D. Technical pentabromodiphenyl ether and hexabromocyclododecane as activators of the pregnane-X-receptor (PXR). *Toxicology*. 2009;264(1-2):45-51.
- Forrest D., Reh TA, Rüsç A. Neurodevelopmental control by thyroid hormone receptors. *Curr Opi. Neurobiol*. 2002, 12, 49-56
- Fängström B, Athanassiadis I, Odsjö T, Norén K, Bergman A. Temporal trends of polybrominated diphenyl ethers and hexabromocyclododecane in milk from Stockholm mothers, 1980-2004. *Mol Nutr Food Res*. 2008;52(2):187-93.
- Gerecke AC, Giger W, Hartmann PC, Heeb NV, Kohler HP, Schmid P, Zennegg M, Kohler M. Anaerobic degradation of brominated flame retardants in sewage sludge. *Chemosphere*. 2006;64(2):311-7.
- Goosey E, Abdallah M, Harrad S. Dust from Primary School and Nursery Classrooms in the UK: Its Significance as a Pathway of Exposure for Young Children to PFOS, PFOA, HBCDs and TBBP-A. *Organohalogen Compd*. 2008; 70: 855-858.
- Gouin T, Harner T. Modelling the environmental fate of the polybrominated diphenyl ethers. *Environ Int*. 2003;29(6):717-24.
- {Great Lakes Chemical Corporation} Material Safety Data Sheet. Great Lakes CD-75-P, CD-75PM and CD-75PC. West Lafayette, Indiana: Great Lakes Chemical Corporation. MSDS Number: 00177. Effective Date: 10/14/2005. 2005, 7 pp.
- Great Lakes Chemical Corporation. 1994; West Lafayette, IN. USA. As cited in: European Commission, 2008
- Guerra P, Martínez E, Farré M, Eljarrat E, Barceló D. Hexabromocyclododecane in human breast milk: Levels and enantiomeric patterns. *Organohalogen Compd*. 2008a, 70, 000309- 000312.
- Hamers T, Kamstra JH, Sonneveld E, Murk AJ, Kester MH, Andersson PL, Legler J, Brouwer A. In vitro profiling of the endocrine-disrupting potency of brominated flame retardants. *Toxicol Sci*. 2006;92(1):157-73.
- Harrad S, Abdallah MA, Covaci A. Causes of variability in concentrations and diastereomer patterns of hexabromocyclododecanes in indoor dust. *Environ Int*. 2009;35(3):573-9.
- Harrad S, Goosey E, Desborough J, Abdallah MA, Roosens L, Covaci A. Dust from U.K. primary school classrooms and daycare centers: the significance of dust as a pathway of exposure of young U.K. children to brominated flame retardants and polychlorinated biphenyls. *Environ Sci Technol*. 2010a;44(11):4198-202.
- Harrad S, de Wit CA, Abdallah MA, Bergh C, Björklund JA, Covaci A, Darnerud PO, de Boer J, Diamond M, Huber S, Leonards P, Mandalakis M, Ostman C, Haug LS, Thomsen C, Webster TF. Indoor contamination with hexabromocyclododecanes, polybrominated diphenyl ethers, and perfluoroalkyl compounds: an important exposure pathway for people? *Environ Sci Technol*. 2010b; 44(9):3221-31.
- Haukås, M, Mariussen, E, Ruus, A, and Tollefsen, KE. Accumulation and disposition of hexabromocyclododecane (HBCD) in juvenile rainbow trout (*Oncorhynchus mykiss*). *Aquatic Toxicology* 2009; 95: 144-151.
- Hayward S, Lei Y and Wania F. Comparative evaluation of three high-performance liquid chromatography-based Kow estimation methods for highly hydrophobic organic compounds: polybrominated diphenyl ethers and hexabromocyclododecane. *Environ Toxicol Chem* 2006; 25: (8): 2018-2027.
- Heeb NV, Schweizer WB, Mattrel P, Haag R, Kohler M, Schmid P, Zennegg M, Wolfensberger M. Regio- and stereoselective isomerization of hexabromocyclododecanes (HBCDs): kinetics and mechanism of beta-HBCD racemization. *Chemosphere*. 2008;71(8):1547-56.
- Heeb NV, Schweizer WB, Kohler M and Gerecke AC. Structure elucidation of hexabromocyclododecanes - a class of compounds with a complex stereochemistry. *Chemosphere* 2005; 61: 65-73.

- Helgason LB, Polder A, Føreid S, Baek K, Lie E, Gabrielsen GW, Barrett RT, Skaare JU. Levels and temporal trends (1983-2003) of polybrominated diphenyl ethers and hexabromocyclododecanes in seabird eggs from north Norway. *Environ Toxicol Chem.* 2009; 28(5):1096-103.
- Hiebl J and Vetter W. (2007) Detection of hexabromocyclododecane and its metabolite pentabromocyclododecane in chicken egg and fish from the official food control. *Journal of Agricultural and Food Chemistry* 2007; 55: 3319-3324.
- Hoh E, Hites RA. 2005. Brominated flame retardants in the atmosphere of the east-central United States. *Environ Sci Technol* 2005; 39(20): 7794-7802.
- Hu J, Liang Y, Chen M, Wang X. Assessing the toxicity of TBBPA and HBCD by zebrafish embryo toxicity assay and biomarker analysis. *Environ Toxicol.* 2009;24(4):334-42.
- [Fraunhofer] 2010, Ruedel H, Mueller J, Ricking M, Quack M, Klein R. Environmental monitoring of HBCD in Europe. Fraunhofer IME, Germany.
- {INE-SEMARNAT} (Instituto Nacional de Ecología, Secretaría de Medio Ambiente y Recursos Naturales). Las sustancias tóxicas persistentes en Mexico. Report. Authors: Bremauntz AF, Yarto Ramírez MY, Díaz JC. 2004, 261 pp.
- {Institut Fresenius} Analysis of hexabromocyclododecane (HBCD) in sludge of the wastewater treatment plant of Broomchemie B.V. in Terneuzen. Taunusstein (DE): Chemische und Biologische Laboratorien GmbH. 2000a.
- {Institut Fresenius} Analysis of hexabromocyclododecane (HBCD) in an effluent of the wastewater treatment plant of Broomchemie B.V. in Terneuzen. Taunusstein (DE): Chemische und Biologische Laboratorien GmbH. 2000b.
- {Institute of Hydrobiology} Validation of the preliminary EU-concept of assessing the impact of chemicals to organisms in sediment by using selected substances. UBA-FB 299 67 411, pp 97 pp. Dresden University of Technology, Dresden, Germany. Authors: Oetken M, Ludwichowski K-U and Nagel R. 2001
- {IPEN}. Next generation of POPs: PBDEs and lindane, International POPs Elimination Network (2005) <http://www.ipen.org/ipenweb/documents/work%20documents/the%20new%20generation%20of%20pops.pdf> (accessed on June 14, 2010)
- Isobe T, Ramu K, Kajiwara N, Takahashi S, Lam PKS, Jefferson TA, Zhou K, Tanabe S. Isomer specific determination of hexabromocyclododecanes (HBCDs) in small cetaceans from the South China Sea – levels and temporal variation. *Marine Pollut Bull* 2008;54: 1139-1145.
- Janák K, Sellström U, Johansson AK, Becher G, de Wit CA, Lindberg P, Helander B. Enantiomer-specific accumulation of hexabromocyclododecanes in eggs of predatory birds. *Chemosphere.* 2008;73(1 Suppl):S193-200.
- Janák K, Covaci A, Voorspoels S, Becher G. Hexabromocyclododecane in marine species from the Western Scheldt estuary: diastereoisomer- and enantiomer-specific accumulation. *Environ Sci Technol* 2005;39(7):1987-1994.
- Jenssen BM, Sørmo EG, Baek K, Bytingsvik J, Gaustad H, Ruus A, Skaare JU. Brominated flame retardants in North-East Atlantic marine ecosystems. *Environ Health Perspect.* 2007;115 Suppl 1:35-41.
- Johnson-Restrepo B, Adams DH, Kannan K. Tetrabromobisphenol A (TBBPA) and hexabromocyclododecanes (HBCDs) in tissues of humans, dolphins, and sharks from the United States. *Chemosphere* 2008;70: 1935-1944.
- Kajiwara N, Sueoka M, Ohiwa T, Takigami H. Determination of flame-retardant hexabromocyclododecane diastereomers in textiles. *Chemosphere.* 2009;74(11):1485-9.
- Kajiwara N, Kamikawa S, Ramu K, Ueno D, Yamada TK, Subramanian A, Lam PK, Jefferson TA, Prudente M, Chung KH, Tanabe S. Geographical distribution of polybrominated diphenyl ethers (PBDEs) and organochlorines in small cetaceans from Asian waters. *Chemosphere.* 2006;64(2):287-95.
- Kakimoto K, Akutsu K, Konishi Y, Tanaka Y. Time trend of hexabromocyclododecane in the breast milk of Japanese women. *Chemosphere.* 2008;71(6):1110-4.
- {KEMI} –(Swedish Chemicals Agency). Proposal for Harmonised Classification and Labelling Based on the CLP Regulation (EC) No 1272/2008, Annex VI, Part 2. Substance Name: Hexabromocyclododecan. Dossier submitted to the European Commission 2009; 49 pp.
- {KLIF} (Climate and Pollution Agency, Norway). Temporal trends of brominated flameretardants, cyclododeca-1,5,9-triene and mercurin in eggs of four seabird species from Northern Norway and Svalbard. Norwegian Pollution Control Authority. Report 942/2005, Authors: Knudsen LB, Gabrielsen GW, Verreault J, Barrett R, Skåre JU, Polder A, Lie E 2005, 44 pp.

- {KLIF} (Climate and Pollution Agency, Norway). Halogenated organic contaminants and mercury in dead or dying seabirds on Bjørnøya (Svalbard). Report 997/2007, TA-number 2222/2007. Authors: Knudsen LB, Sagerup K, Polder A, Slabach M, Josefsen TD, Strøm, Skaare, JU, Gabrielsen GW. 2007, 45 pp.
- {KLIF} (Climate and Pollution Agency, Norway). New organic pollutants in air, 2007. Brominated flame retardants and polyfluorinated substances. SPFO-report 1077/2010, TA-2689/2010. Conducted by the Norwegian Institute for Air Research (NILU). Authors: Manø S, Herzke D, Schlabach M. 2010. 64 pp.  
<http://www.klif.no/no/Tema/Miljoovervakning/Statlig-miljoovervakning/Kartlegging-av-nye-miljogifter/Rapporter/Nye-miljogifter-i-luft-2007/>.
- Kling P, Förlin L. Proteomic studies in zebrafish liver cells exposed to the brominated flame retardants HBCD and TBBPA. *Ecotoxicol Environ Saf.* 2009;72(7):1985-93.
- Knutsen HK, Kvalem HE, Thomsen C, Frøshaug M, Haugen M, Becher G, Alexander J, Meltzer HM. Dietary exposure to brominated flame retardants correlates with male blood levels in a selected group of Norwegians with a wide range of seafood consumption. *Mol Nutr Food Res.* 2008;52(2):217-27.
- Knudsen LB, Borgå, Jørgensen EH, van Bavel B, Schlabach M, Verreault J, Gabrielsen GW. Halogenated organic contaminants and mercury in northern fulmars (*Fulmarus glacialis*): levels, relationships to dietary descriptors and blood to liver comparison. *Environ Pollut* 2007;146: 25-33.
- Kohler M, Zennegg M, Bogdal C, Gerecke AC, Schmid PP, Heeb NV, Sturm M, Vonmont H, Kohler HPE, Giger W.. Temporal trends, congener patterns, and sources of octa-, nona-, and decabromodiphenyl ethers (PBDEs) and hexabromocyclododecanes (HBCD) in Swiss lake sediments. *Environ Sci Technol* 2008;42: 6378-6384.
- Kuiper RV, Canton RF, Leonards PEG, Jenssen BM, Dubbeldam M, Wester PW, van den Berg M, Vos JG, Vethaak AD. Long-term exposure of European flounder (*Platichthys flesus*) to the flame-retardants tetrabromobisphenol A (TBBPA) and hexabromocyclododecane (HBCD). *Ecotox. Environ. Saf.* 2007;67:349-360.
- Kupper T, de Alencastro LF, Gatsigazi R, Furrer R, Grandjean D, Tarradellas J. Concentrations and specific loads of brominated flame retardants in sewage sludge. *Chemosphere.* 2008;71(6):1173-80.
- Larsen, E. R.; Ecker, E. L. Thermal stability of fire retardants: I, Hexabromocyclododecane. *J. Fire Sci.* 1986, 4, 261-275.
- Law RJ, Bersuder P, Barry J, Wilford BH, Allchin CR, Jepson PD. A significant downturn in levels of hexabromocyclododecane in the blubber of harbour porpoises (*Phocoena phocoena*) stranded or bycaught in the UK: an update to 2006. *Environ Sci Technol* 2008a;42: 9104-9109.
- Law RJ, Herzke D, Harrad S, Morris S, Bersuder P, Allchin CR. Levels and trends of HBCD and BDEs in the European and Asian environments, with some information for other BFRs. *Chemosphere.* 2008b;73(2):223-41.
- Law K, Halldorson T, Danell R, Stern G, Gewurtz S, Alaee M, Marvin C, Whittle M, Tomy G. Bioaccumulation and trophic transfer of some brominated flame retardants in a Lake Winnipeg (Canada) food web. *Environ Toxicol Chem.* 2006a;25(8):2177-86. (Erratum in: *Environ Toxicol Chem.* 2007 Jan;26(1):190).
- Law RJ, Allchin CR, de Boer J, Covaci A, Herzke D, Lepom P, Morris S, de Wit CA. Levels and trends of brominated flame retardants in the European environment. *Chemosphere* 2006b: 64: 187-208.
- Law K, Palace VP, Halldorson T, Danell R, Wautier K, Evans B, Alaee M, Marvin C, Tomy GT. Dietary accumulation of hexabromocyclododecane diastereoisomers in juvenile rainbow trout (*Oncorhynchus mykiss*) I: bioaccumulation parameters and evidence of bioisomerization. *Environ Toxicol Chem.* 2006c;25(7):1757-61.
- Law RJ, Bersuder P, Allchin CR, Barry J. Levels of the flame retardants hexabromocyclododecane and tetrabromobisphenol A in the blubber of harbour porpoises (*Phocoena phocoena*) stranded or bycaught in the U.K., with evidence for an increase in HBCD concentrations in recent years. *Environ Sci Technol* 2006d;40: 2177-2183.
- Law RJ, Kohler M, Heeb NV, Gerecke AC, Schmid P, Voorspols S, Covaci A, Becher G, Janak K, and C Thomsen. Hexabromocyclododecane challenges scientists and regulators. *Environ Sci Technol* 2005, 39(13): 281A-287A.
- Leonards P, Vethaak D, Brandsma S, Kwadijk C, Micic D, Jol J, Schout P, de Boer J. Species-specific accumulation and biotransformation of polybrominated diphenyl ethers and hexabromocyclododecane in two Dutch food chains. Abstract. The Third International Workshop on Brominated Flame Retardants, June 6-9, Toronto, Canada. 2004;283-286.
- Letcher RJ, Gebbink WA, Sonne C, Born EW, McKinney MA, Dietz R. Bioaccumulation and biotransformation of brominated and chlorinated contaminants and their metabolites in ringed seals (*Pusa hispida*) and polar bears (*Ursus maritimus*) from East Greenland. *Environ Int.* 2009;35(8):1118-24.

- Lignell S, Aune M, Darnerud PO, Cnattingius S, Glynn A. Persistent organochlorine and organobromine compounds in mother's milk from Sweden 1996-2006: compound-specific temporal trends. *Environ Res.* 2009;109(6):760-7.
- Lilienthal H, van der Ven LT, Piersma AH, Vos JG. Effects of the brominated flame retardant hexabromocyclododecane (HBCD) on dopamine-dependent behavior and brainstem auditory evoked potentials in a one-generation reproduction study in Wistar rats. *Toxicol Lett.* 2009;185(1):63-72.
- Lindberg P, Sellström U, Hågberg L, de Wit CA. Higher brominated diphenyl ethers and hexabromocyclododecane found in eggs of peregrine falcons (*Falco peregrinus*) breeding in Sweden. *Environ Sci Technol* 2004;38: 93-96.
- López D, Athasiadou M, Athanassiadis I, Estrada LY, Díaz-Barriga F and Bergman Å. A preliminary study on PBDEs and HBCDD in blood and milk from Mexican women. In *The third international workshop on brominated flame retardants - BFR 2004*. Book of abstracts. Edited by Alae M and et al. 2004; 483-487.
- Lower N, Moore A. The impact of a brominated flame retardant on smoltification and olfactory function in Atlantic salmon (*Salmo salar* L.) smolts. *Mar. Freshw. Behav. Physiol.* 2007;40:267-284.
- {LCSP} (Lowell Center For Sustainable Production). An Overview of Alternatives to Tetrabromobisphenol A (TBBPA) and Hexabromocyclododecane (HBCD). Report prepared for The Jennifer Altman Foundation. University of Massachusetts, 2006. Author: Morose G. 32 pp.
- Maden M. Retinoic acid in the development, regeneration and maintenance of the nervous system. *Nat. Rev. Neurosci.* 2007, 8, 755-765
- MacGregor JA and Nixon WB. Determination of water solubility of hexabromocyclododecane (HBCD) using a generator column method. 2004; pp 52. Wildlife International, Ltd., Easton, Maryland, USA.
- MacGregor JA and Nixon WB. Hexabromocyclododecane (HBCD): Determination of n-octanol/ water partition coefficient. 1997; 439C- 104, pp 68 pp. Wildlife International Ltd, Easton, Maryland, USA.
- Malarvannan G, Kunisue T, Isobe T, Sudaryanto A, Takahashi S, Prudente M, Subramanian A, Tanabe S. Organohalogen compounds in human breast milk from mothers living in Payatas and Malate, the Philippines: levels, accumulation kinetics and infant health risk. *Environ Pollut.* 2009;157(6):1924-32.
- Managaki S, Miyake Y, Yokoyama Y, Hondo H, Masunaga S, Nakai S, Kobayashi T, Kameya T, Kimura A, Nakarai T, Oka Y, Otani H and Miyake A. Emission load of hexabromocyclododecane in Japan based on the substance flow analysis. 2009. [http://risk.kan.ynu.ac.jp/publish/managaki/managaki200908\\_1.pdf](http://risk.kan.ynu.ac.jp/publish/managaki/managaki200908_1.pdf)
- Mariussen E and Fonnum F. The effect of brominated flame retardants on neurotransmitter uptake into rat brain synaptosomes and vesicles. *Neurochem Int* 2003; 43: 533-542.
- Meaney MJ, Stewart J, Poulin P, McEwen BS. Sexual differentiation of social play in rat pups is mediated by the neonatal androgen-receptor system. *Neuroendocrinology* 1983;37:85-90
- Meijer L, Weiss J, Van Velzen M, Brouwer A, Bergman A, Sauer PJ. Serum concentrations of neutral and phenolic organohalogen compounds in pregnant women and some of their infants in The Netherlands. *Environ Sci Technol.* 2008;42(9):3428-33.
- Miljeteig C, Strøm H, Gavrilov MV, Volkov A, Jenssen BM, Gabrielsen GW. High levels of contaminants in ivory gull *Pagophila eburnea* eggs from the Russian and Norwegian Arctic. *Environ Sci Technol.* 2009 ;43(14):5521-8.
- Minh NH, Isobe T, Ueno D, Matsumoto K, Mine M, Kajiwara N, Takahashi S, Tanabe S. Spatial distribution and vertical profile of polybrominated diphenyl ethers and hexabromocyclododecanes in sediment core from Tokyo Bay, Japan. *Environ Pollut.* 2007;148(2):409-17.
- Miyake Y, Managaki S, Yokoyama Y, Nakai S, Kataoka T, Nagasawa E, Shimojima M, Masunaga S, Hondo H, Kobayashi T, Kameya T, Kimura A, Nakarai T, Oka Y, Otani H and Miyake A. Exposure to hexabromocyclododecane (HBCD) emitted into indoor air by drawing flameretarded curtain. For online access go to: [risk.kan.ynu.ac.jp/publish/masunaga/masunaga200908\\_3.pdf](http://risk.kan.ynu.ac.jp/publish/masunaga/masunaga200908_3.pdf)
- {MOEJ} (Ministry of the Environment, Japan). 6-Week Administration Study of 1,2,5,6,9,10-Hexabromocyclododecane for Avian Reproduction Toxicity under Long-day Conditions using Japanese Quail. Report. Research Institute for Animal Science in Biochemistry & Toxicology, 2009
- Montzka SA, Spivakovsky CM, Butler JH, Elkins JW, Lock LT, Mondeel DJ. New observational constraints for atmospheric hydroxyl on global and hemispheric scales. *Science.* 2000;288(5465):500-3.

- Morf L, Buser A, Taverna R, Bader HP, Scheidegger R. Dynamic substance flow analysis as a valuable tool - a case study for brominated flame retardants as an example of potential endocrine disruptors. 2008; 62(5):424-431
- Morris S, Allchin CR, Zegers BN, Haftka JJ, Boon JP, Belpaire C, Leonards PE, Van Leeuwen SP, de Boer J. Distribution and fate of HBCD and TBBPA brominated flame retardants in North Sea estuaries and aquatic food webs. *Environ Sci Technol*. 2004;38(21):5497-504.
- Muir DCG, Backus S, Derocher AE, Dietz R, Evans TJ, Gabrielsen GW, Nagy J, Norstrom RJ, Sonne C, Stirling I, Taylor MK, Letcher RJ. Brominated flame retardants in polar bears (*Ursus maritimus*) from Alaska, the Canadian Arctic, East Greenland, and Svalbard. *Environ Sci Technol* 2006;40(2): 449-455.
- Muir, D.C.G., Alae, M., Butt, C., Braune, B., Helm, P., Mabury, S., Tomy, G., Wang, X., New contaminants in Arctic biota. In: Synopsis of Research conducted under the 2003–2004, Northern Contaminants Program, Indian and Northern Affairs Canada, Ottawa, ON, Canada, 2004. pp. 139–148.
- {NCM} (Nordic Council of Ministers). Hexabromocyclododecane as a possible global POP. Nordic Chemicals Group and Nordic Council of Ministers, Author: Peltola-Thies J. 2008, 91 pp.
- Norris D. *Vertebrate Endocrinology* 4<sup>th</sup> edition. Elsevier Academic Press, London, UK. 2007. 560 pp, ISBN-13: 978-0-12-088768-2
- Nyholm JR, Norman A, Norrgren L, Haglund P, Andersson PL. Maternal transfer of brominated flame retardants in zebrafish (*Danio rerio*). *Chemosphere*. 2008;73(2):203-8.
- {OECD} (Organization for Economic Co-operation and Development). SIDS Initial Assessment Profile for Cas. No. 25637-99-4, 3194-55-6, Hexabromocyclododecane (HBCDD). SIAM 24, 19-20 April 2007. Available from: <http://webnet.oecd.org/Hpv/UI/handler.axd?id=ea58ac11-e090-4b24-b281-200ae351686c>
- Oppenheim RW. Cell death during development of the nervous system. *Annu Rev Neurosci*. 1991;14:453-501.
- Palace VP, Pleskach K, Halldorson T, Danell R, Wautier K, Evans B, Alae M, Marvin C, Tomy GT. Biotransformation enzymes and thyroid axis disruption in juvenile rainbow trout (*Oncorhynchus mykiss*) exposed to hexabromocyclododecane diastereoisomers. *Environ Sci Technol*. 2008;42(6):1967-72.
- Palace VVP, Park B, Pleskach K, Gemmill B, Tomy G. Altered thyroxine metabolism in rainbow trout (*Oncorhynchus mykiss*) exposed to hexabromocyclododecane (HBCD). *Chemosphere*. 2010 Jun;80(2):165-9.
- Peck AM, Pugh RS, Moors A, Ellisor MB, Porter BJ, Becker PR, Kucklick JR. Hexabromocyclododecane in white-sided dolphins: temporal trend and stereoisomer distribution in tissues. *Environ Sci Technol* 2008;42: 2650-2655.
- Peled M, Scharia R, Sondack D. Thermal rearrangement of hexabromocyclododecane (HBCD). *Adv Organobrom Chem II* 1995; 7: 92-99.
- Petersen M, Hamm S, Schäfer A, Esser U. Comparative GC/MS and LC/MS detection of hexabromocyclododecane (HBCD) in soil and water samples. *Organohalogen Compounds* 66:2005, 226-233
- Polder A, Thomsen C, Lindström G, Løken KB, Skaare JU. Levels and temporal trends of chlorinated pesticides, polychlorinated biphenyls and brominated flame retardants in individual human breast milk samples from Northern and Southern Norway. *Chemosphere*. 2008a;73(1):14-23.
- Polder A, Gabrielsen GW, Odland JØ, Savinova TN, Tkachev A, Løken KB, Skaare JU. Spatial and temporal changes of chlorinated pesticides, PCBs, dioxins (PCDDs/PCDFs) and brominated flame retardants in human breast milk from Northern Russia. *Sci Total Environ*. 2008b;391(1):41-54.
- Polder A, Venter B, Skaare JU, Bouwman M. Polybrominated diphenyl ethers and HBCD in bird eggs of South Africa. *Chemosphere* 2008c;73: 148-154
- {Polymer Research Centre} Final report on flame retardant release from textiles. University of Surrey, UK. 2006. Authors: Thomas J and Stevens G.
- Prinn RG, Weiss RF, Miller BR, Huang J, Alyea FN, Cunnold DM, Fraser PJ, Hartley DE, Simmonds PG. Atmospheric Trends and Lifetime of CH<sub>3</sub>CCl<sub>3</sub> and Global OH Concentrations. *Science*. 1995;269(5221):187-192.
- Pulkrabová J, Hrádková P, Hajslová J, Poustka J, Nápravníková M, Poláček V. Brominated flame retardants and other organochlorine pollutants in human adipose tissue samples from the Czech Republic. *Environ Int*. 2009;35(1):63-8.
- Remberger M, Sternbeck J, Palm A, Kaj L, Strömberg K, Brorström-Lundén E. The environmental occurrence of hexabromocyclododecane in Sweden. *Chemosphere*. 2004;54(1):9-21.



- Reistad T, Fonnum F, Mariussen E. Neurotoxicity of the pentabrominated diphenyl ether mixture, DE-71, and hexabromocyclododecane (HBCD) in rat cerebellar granule cells in vitro. *Arch Toxicol.* 2006;80(11):785-96.
- Roosens L, Cornelis C, D'Hollander W, Bervoets L, Reynders H, Van Campenhout K, Van Den Heuvel R, Neels H, Covaci A. Exposure of the Flemish population to brominated flame retardants: model and risk assessment. *Environ Int.* 2010;36(4):368-76.
- Roosens L, Abdallah MA, Harrad S, Neels H, Covaci A. Exposure to hexabromocyclododecanes (HBCDs) via dust ingestion, but not diet, correlates with concentrations in human serum: preliminary results. *Environ Health Perspect.* 2009;117(11):1707-12.
- Roze E, Meijer L, Bakker A, Van Braeckel KN, Sauer PJ, Bos AF. Prenatal exposure to organohalogenes, including brominated flame retardants, influences motor, cognitive, and behavioral performance at school age. *Environ Health Perspect.* 2009 ;117(12):1953-8.
- Saegusa Y, Fujimoto H, Woo GH, Inoue K, Takahashi M, Mitsumori K, Hirose M, Nishikawa A, Shibutani M. Developmental toxicity of brominated flame retardants, tetrabromobisphenol A and 1,2,5,6,9,10 hexabromocyclododecane, in rat offspring after maternal exposure from mid-gestation through lactation. *Reprod Toxicol.* 2009;28(4):456-67.
- Schantz SL, Widholm JJ. Cognitive effects of endocrine-disrupting chemicals in animals. *Environ Health Perspect.* 2001;109(12):1197-206.
- Schechter A, Harris TR, Shah N, Musumba A, Pöpke O. Brominated flame retardants in US food. *Mol Nutr Food Res.* 2008;52(2):266-72.
- Schriks M, Zvinavashe E, Furlow JD, Murk AJ. Disruption of thyroid hormone-mediated *Xenopus laevis* tadpole tail tip regression by hexabromocyclododecane (HBCD) and 2,2',3,3',4,4',5,5',6-nona brominated diphenyl ether (BDE206). *Chemosphere.* 2006;65(10):1904-8.
- Sellström U, Kierkegaard A, Alsberg T, Jonsson P, Wahlberg C, de Wit C. Brominated flame retardants in sediments from European estuaries, the Baltic Sea and in sewage sludge. *Organohalogen Compd* 1999a;40: 383-386
- Sellström U. Determination of some polybrominated flame retardants in biota, sediment and sewage sludge. Doctoral Dissertation. Stockholm (SE): Stockholm University. 1999b, 71 pp.
- Sellström U, Kierkegaard A, de Wit C, Jansson B. Polybrominated diphenyl ethers and hexabromocyclododecane in sediment and fish from a Swedish river. *Environ Toxicol Chem* 1998;17(6):1065-1072.
- SETAC 2009, Letcher RJ, Fernie KJ, Mattioli LC, Chu SG, Bird D, Ritchie IJ, Schutt LJ. Uptake, Depuration and Fate of Hexabromocyclododecane (HBCD) and BDE-209 Flame Retardants in Dietary Exposed American kestrels (*Falco sparverius*). 30<sup>th</sup> Annual Meeting of SETAC, November 19-23, New Orleans, L.A., U.S.A.
- Shi ZX, Wu YN, Li JG, Zhao YF, Feng JF. Dietary exposure assessment of Chinese adults and nursing infants to tetrabromobisphenol-A and hexabromocyclododecanes: occurrence measurements in foods and human milk. *Environ Sci Technol.* 2009;43(12):4314-9.
- Simpson ER, Clyne C, Rubin G, Boon WC, Robertson K, Britt K, Speed C, Jones M. Aromatase-a brief review. *Annu Rev Physiol* 2002;64:93-127.
- Smith K, Liu C-H, El-Hiti GA, Kang GS, Jonas E, Clement SG, Checquer AD, Howarth OW, Hursthouse MB and Coles SJ. An extensive study of bromination of cis, trans-1,5,9-cyclododecatriene: product structures and conformations. *Org Biomol Chem* 2005; 3: 1880-1892.
- Stapleton HM, Allen JG, Kelly SM, Konstantinov A, Klosterhaus S, Watkins D, McClean MD, Webster TF. Alternate and new brominated flame retardants detected in U.S. house dust. *Environ Sci Technol.* 2008;42(18):6910-6.
- Stapleton HM, Dodder NG, Kucklick JR, Reddy CM, Schantz MM, Becker PR, Gulland F, Porter BJ, Wise SA. Determination of HBCD, PBDEs and MeO-BDEs in California sea lions (*Zalophus californianus*) stranded between 1993 and 2003. *Mar Pollut Bull* 2006;52: 522-531.
- Stenzel JI and Nixon WB. Hexabromocyclododecane (HBC): Determination of the vapor pressure using a spinning rotor gauge. 1997; 439C-117, pp 44. Wildlife international Ltd., Easton, Maryland.
- Stenzel JI and Markley BJ. Hexabromocyclododecane (HBCD): Determination of the water solubility. 1997; 439C-105, 55 pp. Wildlife International Ltd, Easton, Maryland, USA.

- Stuart H, Ibarra C, Abdallah MA, Boon R, Neels H, Covaci A. Concentrations of brominated flame retardants in dust from United Kingdom cars, homes, and offices: Causes of variability and implications for human exposure. *Environ Int.* 2008;34(8):1170-5.
- {Swerea} Exploration of management options for HBCDD. Report. Authors: Posner S, Roos S, Olsson E. 2010. 84 pp.
- Sørmo EG, Salmer MP, Jenssen BM, Hop H, Baek K, Kovacs KM, Lydersen C, Falk-Petersen S, Gabrielsen GW, Lie E, Skaare JU. Biomagnification of polybrominated diphenyl ether and hexabromocyclododecane flame retardants in the polar bear food chain in Svalbard, Norway. *Environ Toxicol Chem.* 2006;25(9):2502-11.
- Takigami H, Suzuki G, Hirai Y, Ishikawa Y, Sunami M, Sakai S. Flame retardants in indoor dust and air of a hotel in Japan. *Environ Int.* 2009a;35(4):688-93.
- Takigami H, Suzuki G, Hirai Y, Sakai S. Brominated flame retardants and other polyhalogenated compounds in indoor air and dust from two houses in Japan. *Chemosphere.* 2009b ;76(2):270-7.
- Tanabe S. Temporal trends of brominated flame retardants in coastal waters of Japan and South China: retrospective monitoring study using archived samples from es-Bank, Ehime University, Japan. *Mar Pollut Bull* 2008;57: 267-274.
- Thomsen C, Knutsen HK, Liane VH, Frøshaug M, Kvalem HE, Haugen M, Meltzer HM, Alexander J, Becher G. Consumption of fish from a contaminated lake strongly affects the concentrations of polybrominated diphenyl ethers and hexabromocyclododecane in serum. *Mol Nutr Food Res.* 2008;52(2):228-37.
- Thomsen C, Molander P, Daae HL, Janák K, Frøshaug M, Liane VH, Thorud S, Becher G, Dybing E. Occupational exposure to hexabromocyclododecane at an industrial plant. *Environ Sci Technol.* 2007;41 (15):5210-6.
- Tomy GT, Pleskach K, Ferguson SH, Hare J, Stern G, Macinnis G, Marvin CH, Loseto L. Trophodynamics of some PFCs and BFRs in a western Canadian Arctic marine food web. *Environ Sci Technol.* 2009;43(11):4076-81.
- Tomy, GT, Pleskach, K, Oswald, T, Halldorson, THJ, Helm, PA, Marvin, CH, MacInnis, G. Enantioselective bioaccumulation of hexabromocyclododecane and congener-specific accumulation of brominated diphenyl ethers in an Eastern Canadian Arctic marine food web. *Environ Sci Technol* 2008: 42: 3634-3639.
- Tomy GT, Budakowski WR, Halldorson THJ, Whittle DM, Keir MJ, MacInnis G, Alae M. Biomagnification of  $\alpha$ - and  $\delta$ -hexabromocyclododecane (HBCD) isomers in a Lake Ontario food web. *Environ Sci Technol* 2004a;38: 2298-2303.
- Tomy GT, Halldorson T, Danell R, Law K, Stern G, Gerwutz S, Whittle M, Alae M, Marvin C. Hexabromocyclododecane (HBCD) isomers and brominated diphenyl ether (BDE) congeners in fish from Lake Winnipeg, Manitoba (Canada). Abstract. Third International Workshop on Brominated Flame Retardants, June 6-9, Toronto, Canada. 2004b. 213-216.
- Tue NM, Sudaryanto A, Minh TB, Isobe T, Takahashi S, Viet PH, Tanabe S. Accumulation of polychlorinated biphenyls and brominated flame retardants in breast milk from women living in Vietnamese e-waste recycling sites. *Sci Total Environ.* 2010;408(9):2155-62.
- Ueno D, Alae M, Marvin C, Muir DCG, Macinnis G, Reiner E, Crozier P, Furdui VI, Subramanian A, Fillmann G, Lam PKS, Zheng GJ, Muchtar M, Razak H, Prudente M, Chung K, Tanabe S. Distribution and transportability of hexabromocyclododecane (HBCD) in the Asia-Pacific region using skipjack tuna as a bioindicator. *Environ Pollut* 2006;144: 238-247.
- {US EPA} {US Environmental Protection Agency}. Initial Risk-Based Prioritization of High Production Volume Chemicals. Chemical/Category: Hexabromocyclododecane (HBCD). Risk-Based Prioritization Document 3/18/2008
- Veith GD, DeFoe DL, and Bergstedt BV. Measuring and estimating the bioconcentration factor of chemicals in fish. *Journal of the Fisheries Research Board of Canada* 1979;36: 1040-1048.
- van der Ven LT, van de Kuil T, Leonards PE, Slob W, Lilienthal H, Litens S, Herlin M, Håkansson H, Cantón RF, van den Berg M, Visser TJ, van Loveren H, Vos JG, Piersma AH. Endocrine effects of hexabromocyclododecane (HBCD) in a one-generation reproduction study in Wistar rats. *Toxicol Lett.* 2009;185(1):51-62.
- van der Ven LT, Verhoef A, van de Kuil T, Slob W, Leonards PE, Visser TJ, Hamers T, Herlin M, Håkansson H, Olausson H, Piersma AH, Vos JG. A 28-day oral dose toxicity study enhanced to detect endocrine effects of hexabromocyclododecane in Wistar rats. *Toxicol Sci.* 2006;94(2):281-92.
- {Velsicol Chemicals} Pharmacokinetics of HBCD in rats. unpublished paper translated into English. Authors: Yu, CC, Atallah, YH. 1980.

- Verreault J, Shahmiri S, Gabrielsen GW, Letcher RJ. Organohalogen and metabolically-derived contaminants and associations with whole body constituents in Norwegian Arctic glaucous gulls. *Environ Int* 2007. 33: 823-830.
- Walsh GE, Yoder MJ, McLaughlin LL. Responses of Marine Unicellular Algae to Brominated Organic Compounds in Six Growth Media. *Ecotoxicology and Environmental Safety* 1987;14, 215-222.
- Wania F, Dugani CB. 2003. Assessing the long-range transport potential of polybrominated diphenyl ethers: a comparison of four multimedia models. *Environ Toxicol Chem* 22(6): 1252-1261.
- Weiss J, Wallin E, Axmon A, Jönsson BA, Akesson H, Janák K, Hagmar L, Bergman A. Hydroxy-PCBs, PBDEs, and HBCDDs in serum from an elderly population of Swedish fishermen's wives and associations with bone density. *Environ Sci Technol*. 2006;40(20):6282-9.
- Weiss J, Meijer L, Sauer P, Linderholm L, Athanassiadis I, Bergman, Å.. PBDE and HBCDD levels in blood from Dutch mothers and infants- Analysis of a Dutch Groningen Infant Cohort. *Organohalogen Compd* 2004;66: 2677-2682.
- {WIL}\_ (WIL Research Laboratories). A 90-day oral (gavage) toxicity study of HBCD in rats. WIL-186012, pp 1527. WIL Research Laboratories, Inc, Ashland, Ohio, USA. Author: Chengelis CP. 2001.
- {Wildlife International}\_ (Wildlife International; Ltd, Easton, Maryland, USA). Hexabromocyclododecane (HBCD): Closed bottle test. 439E-102. 1996. Authors: Schaefer E. and Haberlein D.
- {Wildlife International}\_ (Wildlife International; Ltd, Easton, Maryland, USA). Hexabromocyclododecane (HBCD): a 48-hour flow-through acute toxicity test with the cladoceran (*Daphnia magna*). Project Number: 439A-102. Wildlife. Authors: Graves W, Swigert J. 1997
- {Wildlife International}\_ (Wildlife International; Ltd, Easton, Maryland, USA). Hexabromocyclododecane (HBCD): A flow-through life-cycle toxicity test with the cladoceran (*Daphnia magna*). Final report. Authors: Drottar KR and Krueger HO. 1998; 439A-108, 78 pp.
- {Wildlife International}\_ (Wildlife International; Ltd, Easton, Maryland, USA). Hexabromocyclododecane (HBCD): A flow-through bioconcentration test with the rainbow trout (*Oncorhynchus mykiss*). Report 439A-108, 1-78. Authors: Drottar KR and Krueger HO. 2000.
- {Wildlife International}\_ (Wildlife International; Ltd, Easton, Maryland, USA). Hexabromocyclododecane (HBCD): A toxicity test to determine the effects of the test substance on seedling emergence of six species of plants. Final report. 2002; 126 pp. Authors: Porch JR, Kendall TZ and Krueger HO.
- {Wildlife International}\_ (Wildlife International; Ltd, Easton, Maryland, USA). Hexabromocyclododecane (HBCD): A 72-hour toxicity test with the marine diatom (*Skeletonema costatum*), Final report. 2004a; Authors: Desjardins D, MacGregor JA and Krueger HO. 66 pp.
- {Wildlife International}\_ (Wildlife International; Ltd, Easton, Maryland, USA). Determination of water solubility of hexabromocyclododecane (HBCD) using a generator column method. 2004b; Authors: MacGregor JA and Nixon WB. pp 52.
- {Wildlife International}\_ (Wildlife International; Ltd, Easton, Maryland, USA). Hexabromocyclododecane (HBCD): A 72-hour toxicity test with the marine diatom (*Skeletonema costatum*) using generator column saturated media. Chapter 2. Final report. 2005; Authors: Desjardins D, MacGregor JA and Krueger HO. 19 pp.
- de Wit CA, Herzke D, Vorkamp K. Brominated flame retardants in the Arctic environment - trends and new candidates. *Sci Total Environ*. 2009. PubMed PMID: 19815253.
- de Wit CA, Alaee M, Muir DC. Levels and trends of brominated flame retardants in the Arctic. *Chemosphere*. 2006;64(2):209-33.
- {WSKOWWIN}\_ Water Solubility for Organic Compounds Program for Microsoft Windows {Estimation Model}\_, Version 1.41. Washington (DC): U.S. Environmental Protection Agency, Office of Pollution Prevention and Toxics; Syracuse (NY): Syracuse Research Corporation. 2000. Available from: [www.epa.gov/oppt/exposure/pubs/episuite.htm](http://www.epa.gov/oppt/exposure/pubs/episuite.htm)
- Xian Q, Ramu K, Isobe T, Sudaryanto A, Liu X, Gao Z, Takahashi S, Yu H, Tanabe S. Levels and body distribution of polybrominated diphenyl ethers (PBDEs) and hexabromocyclododecanes (HBCDs) in freshwater fishes from the Yangtze River, China. *Chemosphere* 2008;71: 268-276.
- Xiao, H., Hung, H., Shen, L., Wania, F., Sverko, E., Su, Y. et al. The atmospheric concentrations of brominated flame retardants at extreme remote locations: The Canadian high Arctic and the Tibetan Plateau. Conference paper.

- Abstract. Fifth International Symposium on Brominated Flame retardants [BFR 2010]. 2010. Kyoto, Japan. 7—10 April 2010.
- Yamada-Okabe T, Sakai H, Kashima Y, Yamada-Okabe H. Modulation at a cellular level of the thyroid hormone receptor-mediated gene expression by 1,2,5,6,9,10-hexabromocyclododecane (HBCD), 4,4'-diiodobiphenyl (DIB), and nitrofen (NIP). *Toxicol Lett.* 2005;155(1):127-33.
- Yu Z, Peng P, Sheng G, Fu J. Determination of hexabromocyclododecane diastereoisomers in air and soil by liquid chromatography-electrospray tandem mass spectrometry. *J Chromatogr A* 2008;1190: 74-79.
- Zegers BN, Mets A, Van Bommel R, Minkenberg C, Hamers T, Kamstra JH, Pierce GJ, Boon JP. Levels of hexabromocyclododecane in harbor porpoises and common dolphins from western European seas, with evidence for stereoisomer-specific biotransformation by cytochrome P450. *Environ Sci Technol.* 2005;39(7):2095-100.
- Zhang X, Yang F, Luo C, Wen S, Zhang X, Xu Y. Bioaccumulative characteristics of hexabromocyclododecanes in freshwater species from an electronic waste recycling area in China. *Chemosphere.* 2009;76(11):1572-8.
- Zhang X, Yang F, Zhang X, Xu Y, Liao T, Song S, Wang J. Induction of hepatic enzymes and oxidative stress in Chinese rare minnow (*Gobiocypris rarus*) exposed to waterborne hexabromocyclododecane (HBCDD). *Aquat Toxicol.* 2008;86(1):4-11.
- Zhang X, Yang F, Xu C, Liu W, Wen S, Xu Y. Cytotoxicity evaluation of three pairs of hexabromocyclododecane (HBCD) enantiomers on Hep G2 cell. *Toxicol In Vitro.* 2008;22(6):1520-7.
- Albemarle Corporation. IUCLID. Data set 201-15946; 2005 (<http://www.epa.gov/hpv/pubs/summaries/cyclodod/c13459rr.pdf>).
- Aniagu SO, Williams TD, Chipman JK. Changes in gene expression and assessment of DNA methylation in primary human hepatocytes and HepG2 cells exposed to the environmental contaminants-Hexabromocyclododecane and 17-beta oestradiol. *Toxicology.* 2009;256(3):143-51.
- Aniagu SO, Williams TD, Allen Y, Katsiadaki I, Chipman JK. Global genomic methylation levels in the liver and gonads of the three-spine stickleback (*Gasterosteus aculeatus*) after exposure to hexabromocyclododecane and 17-beta oestradiol. *Environ Int.* 2008;34(3):310-7.
- Asplund L, Bignert A, Nylund K. Comparison of spatial and temporal trends of methoxylated PBDEs, PBDEs, and hexabromocyclododecane in herring along the Swedish coast. *Organohalogen Compd* 2004;66: 3988-3993.
- {BASF} Hexabromocyclododecane: 28-day feeding trials with rats. BASF (not published). Authors: Zeller H and Kirsch P. 1969.
- {BASF} Cytogenetic study in vivo with hexabromocyclododecane in the mouse micronucleus test after two intraperitoneal administrations. Report. Experimental toxicology and ecology, BASF, Germany. Project no 26M0100/004018; 2000.
- Birnbaum, L.S., Staskal, D.F. Brominated flame retardants: cause for concern? *Environ Health Perspect.* 2004, 112, 9–17.
- Brent GA, Larsen PR, Davies TF. Hypothyroidism and thyroiditis. In: Kronenberg: HM, Shlomo M, Polonsky KR, Larsen PR, eds. *Williams Textbook of Endocrinology.* 11th ed. Saunders Elsevier, Philadelphia, 2008: Chapter 12.
- Bustnes JO, Yoccoz NG, Bangjord G, Polder A, Skaare JU. Temporal trends (1986-2004) of organochlorines and brominated flame retardants in tawny owl eggs from northern Europe. *Environ Sci Technol* 2007;41: 8491-8497.
- Bytingsvik J, Gaustad H, Pettersvik Salmer M, Soermo EG, Baek K, Føreid S, Ruus A, Utne Skaare J, Munro Jenssen B. 2004. Spatial and temporal trends of BFRs in Atlantic cod and Polar cod in the North-East Atlantic. *Organohalogen Compd* 66: 3918-3922.
- Cantón RF, Peijnenburg AA, Hoogenboom RL, Piersma AH, van der Ven LT, van den Berg M, Heneweer M. Subacute effects of hexabromocyclododecane (HBCD) on hepatic gene expression profiles in rats. *Toxicol Appl Pharmacol.* 2008 ;231(2):267-72.
- Carpenter DO. Environmental contaminants as risk factors for developing diabetes. *Rev Environ Health.* 2008;23(1):59-74.

- Catalá A. Lipid peroxidation of membrane phospholipids generates hydroxy-alkenals and oxidized phospholipids active in Environmental physiological and/or pathological conditions. *Chem Phys Lipids*. 2009;157(1):1-11.
- Darnerud PO. Toxic effects of brominated flame retardants in man and in wildlife. *Environ Int*. 2003 Sep;29(6):841-53. Review.
- D'Silva K, Fernandes A, Rose M. 2004. Brominated organic micropollutants – igniting the flame retardant issue. *Crit Rev Env Sci Technol* 34: 141-207.
- Eljarrat E, de la Cal A, Raldua D, Duran C, Barcelo D. Brominated flame retardants in *Alburnus alburnus* from Cinca River basin (Spain). *Environ Pollut* 2005;133: 501-508.
- Eljarrat E, de la Cal A, Raldua D, Duran C, Barcelo D. Occurrence and bioavailability of polybrominated diphenyl ethers and hexabromocyclododecane in sediment and fish from the Cinca River, a tributary of the Ebro River (Spain). *Environ Sci Technol* 2004;38(9): 2603-2608.
- {EQC}\_ Equilibrium Criterion Model. Version 2.02. Peterborough (ON): Trent University, Canadian Environmental Modelling Centre. 2003. Available from: <http://www.trentu.ca/academic/aminss/envmodel/models/EQC2.html>
- Evenset A, Christensen GN, Carroll J, Zaborska A, Berger U, Herzke D, Gregor D. Historical trends in persistent organic pollutants and metals recorded in sediment from Lake Ellasjøen, Bjørnøya, Norwegian Arctic. *Environ Pollut* 2007;146: 196-205.
- Feinberg AP, Tycko B. The history of cancer epigenetics. *Nat. Rev. Cancer* 2004;4:143-153.
- Fernie KJ, Shutt JL, Letcher RJ, Ritchie IJ, Bird DM. Environmentally relevant concentrations of DE-71 and HBCD alter eggshell thickness and reproductive success of American kestrels. *Environ Sci Technol*. 2009;43(6):2124-30.
- Franco R, Schoneveld O, Georgakilas AG, Panayiotidis MI. Oxidative stress, DNA methylation and carcinogenesis. *Cancer Lett*. 2008;266(1):6-11.
- Gauthier LT, Hebert CE, Weseloh DVC, Trudeau S, Letcher RJ. Current-use and emerging brominated and chlorinated flame retardant contaminants in the eggs of herring gulls (*Larus argentatus*) from colonial Great Lakes populations. Poster. 8<sup>th</sup> Annual Workshop on Brominated Flame Retardants in the Environment, June 27-29, Toronto, Canada. 2006.
- Gauthier LT, Hebert CE, Weseloh DVC, Letcher RJ. Current-use flame retardants in the eggs of herring gulls (*Larus argentatus*) from the Laurentian Great Lakes. *Environ Sci Technol* 2007;41: 4561-4567.
- Gebbink WA, Sonne C, Dietz R, Kirkegaard M, Born EW, Muir DCG, Letcher RJ. 2008. Target tissue selectivity and burdens of diverse classes of brominated and chlorinated contaminants in polar bears (*Ursus maritimus*) from East Greenland. *Environ Sci Technol* 2008; 42: 752-759.
- Gee JR, Moser VC. Acute postnatal exposure to brominated diphenylether 47 delays neuromotor ontogeny and alters motor activity in mice. *Neurotoxicol Teratol*. 2008;30(2):79-87.
- Gerecke AC, Kohler M, Zennegg M, Schmid P, Heeb NV. Detection of  $\alpha$ -isomer dominated HBCD (hexabromocyclododecane) in Swiss fish at levels comparable to PBDEs (polybrominated diphenyl ethers). *Organohalogen Compd* 2003; 61: 155-158.
- Germer S, Piersma AH, van der Ven L, Kamyschnikow A, Fery Y, Schmitz HJ, Schrenk D. Subacute effects of the brominated flame retardants hexabromocyclododecane and tetrabromobisphenol A on hepatic cytochrome P450 levels in rats. *Toxicology*. 2006;218(2-3):229-36.
- Guerra P, Eljarrat E, Barceló D. Enantiomeric specific determination of hexabromocyclododecane by liquid chromatography-quadrupole linear ion trap mass spectrometry in sediment samples. *J Chromatogr A* 2008b;1203: 81-87.
- Helleday T, Tuominen K-L, Bergman Å and Jenssen D. Brominated flame retardants induce intragenic recombination in mammalian cells. *Mutat Res* 1999; 439: 137-147.
- {HENRYWIN}\_ Henry's Law Constant Program for Microsoft Windows {Estimation Model}\_ Version 3.10. Washington (DC): U.S. Environmental Protection Agency, Office of Pollution Prevention and Toxics; Syracuse (NY): Syracuse Research Corporation. 2000. Available from: [www.epa.gov/oppt/exposure/pubs/episuite.htm](http://www.epa.gov/oppt/exposure/pubs/episuite.htm)
- Hinkson NC, Whalen MM. Hexabromocyclododecane decreases the lytic function and ATP levels of human natural killer cells. *J Appl Toxicol*. 2009;29(8):656-661.
- Hites RA, Hoh E. Brominated flame retardants in the atmosphere of the U.S. Presentation at the Seventh Annual Workshop on Brominated Flame Retardants (BFR) in the Environment, June 13-14, Gaithersburg, MD. 2005, 37 pp.

- {HSDB} (Hazardous Substances Data Bank, {database on the Internet}). Bethesda (MD): National Library of Medicine (US). 1983-. Available from: <http://toxnet.nlm.nih.gov/cgi-bin/sis/htmlgen?HSDB>
- {HYDROWIN}. Hydrolysis Rates Program for Microsoft Windows {Estimation Model}. Version 1.67. Washington (DC): U.S. Environmental Protection Agency, Office of Pollution Prevention and Toxics; Syracuse (NY): Syracuse Research Corporation. 2000. Available from: [www.epa.gov/oppt/exposure/pubs/episuite.htm](http://www.epa.gov/oppt/exposure/pubs/episuite.htm)
- Ismail N, Gewurtz SB, Pleskach K, Whittle DM, Helm PA, Marvin CH, Tomy GT. Brominated and chlorinated flame retardants in Lake Ontario, Canada, lake trout (*Salvelinus namaycush*) between 1979 and 2004 and possible influences of food web changes. *Environ Toxicol Chem* 2009; 28(5): 910–920.
- Jaspers V, Covaci A, Maervoet J, Dauwe T, Voorspoels S, Schepens P, Eens M. 2005. Brominated flame retardants and organochlorine pollutants in eggs of little owls (*Athene noctua*) from Belgium. *Environ Pollut* 136: 81-88.
- Johansson A, Sellström U, Lindberg P, Bignert A, de Wit CA. Polybrominated diphenyl ether congener patterns, hexabromocyclododecane, and brominated biphenyl 153 in eggs of peregrine falcons (*Falco peregrinus*) breeding in Sweden. *Environ Toxicol Chem* 2009;28(1): 9-17.
- Jones OA, Maguire ML, Griffin JL. Environmental pollution and diabetes: a neglected association. *Lancet*. 2008 26;371(9609):287-8.
- Kemmlin S, Hahn O, Jann O. 2003. Emissions of organophosphate and brominated flame retardants from selected consumer products and building materials. *Atmos Environ* 2003;37: 5485-5493.
- Kezele P, Skinner MK. Regulation of ovarian primordial follicle assembly and development by estrogen and progesterone: endocrine model of follicle assembly. *Endocrinology*. 2003;144(8):3329-37.
- Kierkegaard A, Sellström U, Bignert A, Olsson M, Asplund L, Jansson B, de Wit C. Temporal trends of a polybrominated diphenyl ether (PBDE), a methoxylated PBDE, and hexabromocyclododecane (HBCD) in Swedish biota. *Organohalogen Compd* 1999;40: 367-370.
- Kim WR, Park OH, Choi S, Choi SY, Park SK, Lee KJ, Rhyu IJ, Kim H, Lee YK, Kim HT, Oppenheim RW, Sun W. The maintenance of specific aspects of neuronal function and behavior is dependent on programmed cell death of adult-generated neurons in the dentate gyrus. *Eur J Neurosci*. 2009;29(7):1408-21. E
- Kohler M, Bogdal C, Zennegg M, Schmid P, Gerecke AC, Heeb NV, Sturm M, Zwysig A, Kohler H, Hartmann PC, Giger W, Scheringer M, Hungerbühler K. Temporal trends of the brominated flame retardants decaBDE,  $\Sigma$  tri-hepta BDEs and HBCDs in Swiss lake sediments. Poster. 17<sup>th</sup> Annual SETAC Meeting, May 20-24, Porto, PT. 2007.
- {KOWWIN}. Octanol-Water Partition Coefficient Program for Microsoft Windows {Estimation Model}. Version 1.67. Washington (DC): U.S. Environmental Protection Agency, Office of Pollution Prevention and Toxics; Syracuse (NY): Syracuse Research Corporation. 2000. Available from: [www.epa.gov/oppt/exposure/pubs/episuite.htm](http://www.epa.gov/oppt/exposure/pubs/episuite.htm)
- Kunisue T, Takayanagi N, Isobe T, Takahashi S, Nakatsu S, Tsubota T, Okumoto K, Bushisue S, Tanabe S. Regional trend and tissue distribution of brominated flame retardants and persistent organochlorines in raccoon dogs (*Nyctereutes procyonoides*) from Japan. *Environ Sci Technol* 2008;42: 685-691.
- Larsen R, Davis E, Peck A, Liebert D, Richardson K. Hexabromocyclododecane in Chesapeake Bay fish. Poster. Seventh Annual Workshop on Brominated Flame Retardants (BFRs) in the Environment, June 13-14, Gaithersburg, MD, USA, 2005.
- Law RJ, Bersuder P, Barry J, Wilford BH, Allchin CR, Jepson PD. A significant downturn in levels of hexabromocyclododecane in the blubber of harbour porpoises (*Phocoena phocoena*) stranded or bycaught in the UK: an update to 2006. *Environ Sci Technol* 2008;42: 9104-9109.
- Lee DH, Lee IK, Song KE, Steffes M, Toscano W, Baker BA, et al. A strong dose–response relation between serum concentrations of persistent organic pollutants and diabetes: results from the National Health and Examination Survey. *Diabetes Care* 2006;29:1638–44.
- Lee DH, Steffes M, Jacobs DR. Positive associations of serum concentration of polychlorinated biphenyls or organochlorine pesticides with self-reported arthritis, especially rheumatoid type, in women. *Environ Health Perspect*. 2007;115(6):883-8.
- van Leeuwen S, Traag W, de Boer J. 2004. Monitoring of brominated flame retardants, dioxins, PCBs and other organohalogen compounds in fish from The Netherlands. *Organohalogen Compd* 2004;66: 1764-1769.

- Lema SC, Nevitt GA. Evidence that thyroid hormone induces olfactory cellular proliferation in salmon during a sensitive period for imprinting. *J Exp Biol.* 2004;207(Pt 19):3317-27.
- Lindberg P, Odsjö T. Mercury levels in feathers of peregrine falcon *Falco peregrinus* compared with total mercury content in some of its prey species. *Environ Pollut (Series B)* 1983;5: 297-318
- Litton Bionetics Inc., Kensington, Md. Mutagenicity Evaluation of 421–32B: Final Report. LBI Project No. 2547. Sponsored by Ciba-Geigy Corp., Ardley, New York. EPA/ OTS Doc #86-900000256. Author: Brusick, D. 1976.
- Lowe SW, Lin AW. Apoptosis in cancer. *Carcinogenesis.* 2000;21(3):485-95.
- Lundstedt-Enkel K, Asplund L, Nylund K, Bignert A, Tysklind M, Olsson M, Orberg J. Multivariate data analysis of organochlorines and brominated flame retardants in Baltic Sea guillemot (*Uria aalge*) egg and muscle. *Chemosphere.* 2006;65(9):1591-9.
- Manchester (UK) Report 10531-009-420/PAH-2. Environmental assessment of a European flame retardant coating manufacturing facility (formulator/compounder). Authors: Dames, Moore. 2000a.
- Manchester (UK):. Report 10531-009-420/PAH-1. Environmental assessment of a HBCD European manufacturing plant. Authors: Dames, Moore. 2000a.
- Marsh G, Athanasiadou M, Bergman Å, Athanassiadis I, Endo T, Haraguchi K. Identification of a novel dimethoxylated polybrominated biphenyl bioaccumulating in marine mammals. *Organohalogen Compd* 2004;66: 3823-3829.
- Marvin CH, Tomy GT, Alaee M, MacInnis G. Distribution of hexabromocyclododecane in Detroit River suspended sediments. Abstract. The Third International Workshop on Brominated Flame Retardants, June 6-9, Toronto, Canada. 2004; 137-140.
- Marvin CH, Tomy GT, Alaee M, MacInnis G. Distribution of hexabromocyclododecane in Detroit River suspended sediments. *Chemosphere* 2006;64: 268-275.
- McKinney MA, Cesh LS, Elliott JE, Williams TD, Garcelon DK, Letcher RJ. Brominated flame retardants and halogenated phenolic compounds in North American West Coast bald eaglet (*Haliaeetus leucocephalus*) plasma. *Environ Sci Technol* 2006;40: 6275-6281.
- Microbiological associates, Inc. Rockville, MD. Chromosome aberrations in human peripheral blood lymphocytes. Study no G96A061.342 sponsored by Chemical Manufacturers Association, Arlington, Virginia, USA; 1996. Authors: Gudi R, Scadly EH.
- Morris AD, Muir DCG, Teixeira C., Epp,J., Sturman S, Solomon KR. Bioaccumulation and distribution of brominated flame retardants and current-use pesticides in an Arctic marine food-web. Abstract. Proceedings of Society of Environmental Toxicology and Chemistry. 28th Annual Meeting, Milwaukee, Wisconsin. 2007.
- Müller MJ, Seitz HJ. Thyroid hormone action on interdiciary metabolism. Part II: Lipid metabolism in hypo- and hyperthyroidism. *Klin. Wochenschr.* 2006, 62, 49-55
- Murvoll KM, Skaare JU, Anderssen E, Jenssen BM. Exposure and effects of persistent organic pollutants in European shag (*Phalacrocorax aristotelis*) hatchlings from the coast of Norway. *Environ Toxicol Chem* 2006a;25(1): 190-198.
- Murvoll KM, Skaare JU, Moe B, Anderssen E, Jenssen BM. Spatial trends and associated biological responses of organochlorines and brominated flame retardants in hatchlings of North Atlantic kittiwakes (*Rissa tridactyla*). *Environ Toxicol Chem* 2006b;25(6): 1648-1656.
- Norstrom RJ, Simon M, Moisey J, Wakeford B, Weseloh DVC. Geographical distribution (2000) and temporal trends (1981-2000) of brominated diphenyl ethers in Great Lakes herring gull eggs. *Environ Sci Technol* 2002;36: 4783-4789.
- {OECD} (Organization for Economic Co-operation and Development). Emission scenario document on plastics additives [Internet]. Paris (FR): OECD, Environment Directorate. 2004a. Available from :[http://www.oecd.org/olOKis/2004doc.nsf/LinkTo/NT0000451A/\\$FILE/JT00166678.PDF](http://www.oecd.org/olOKis/2004doc.nsf/LinkTo/NT0000451A/$FILE/JT00166678.PDF).
- {OECD} (Organization for Economic Co-operation and Development). Manual for investigation of HPV chemicals. Paris (FR): OECD Secretariat. April 2003. Available from: [http://www.oecd.org/document/710,2340,en\\_2649\\_34379\\_1947463\\_1\\_1\\_1\\_1.00.html](http://www.oecd.org/document/710,2340,en_2649_34379_1947463_1_1_1_1.00.html).

- OECD (Organization for Economic Co-operation and Development-(~~OECD~~)). Guidelines for the testing of chemicals. Guideline 307. Aerobic and anaerobic transformation in soil. Paris (FR): OECD, Environment Directorate. 2002. 17 pp.
- Parker RM. Testing for reproductive toxicity: In: Hood Rd, Editor. Developmental and reproductive toxicology. 2<sup>nd</sup> edition. CRC Press, Taylor&Francis group, Boca Ration, USA; 2006;435-487.
- {PCK<sub>OC</sub>WIN}\_{ Organic Carbon Partition Coefficient Program for Windows {Estimation Model}\_{, 2000. Version 1.66. Washington (DC): U.S. Environmental Protection Agency, Office of Pollution Prevention and Toxics; Syracuse (NY): Syracuse Research Corporation, USA. Available from: [www.epa.gov/oppt/exposure/pubs/episuite.htm](http://www.epa.gov/oppt/exposure/pubs/episuite.htm)
- Peters RJB, Beeltje H, van Delft RJ. Xeno-estrogenic compounds in precipitation. *J Environ Monit* 2008;10: 760-769.
- Pritchard M, Fournel-Gigleux S, MacKanezie P, Magdalou J. A recombinant Phenobarbital-inducible rat liver UDPglucuronosyltransferase (UDP-glucuronosyltransferase 2B1) stably expressed in V79 cells catalyzes the glucuronidation of morphine, phenols and carboxylic acids. *Mol. Pharmacol.* 1994 ;45: 42–50.
- PWGSC-INAC-NCP. Northern Contaminants Program (NCP) Report II Sources of Occurrence, Trends and Pathways in the Physical Environment. QS-8525-002-EE-A1 2003. Available at <http://www.ainc-inac.c.ca/nth/ct/ncp/pubs.phy/phy-eng.asp>
- Ramu K, Kajiwarra N, Isobe T, Takahashi S, Kim EY, Min BY, We SU, Tanabe S. Spatial distribution and accumulation of brominated flame retardants, polychlorinated biphenyls and organochlorine pesticides in blue mussels (*Mytilus edulis*) from coastal waters of Korea. *Environ Pollut* 2007;148: 562-569.
- Rawn DFK, Sadler A, Lau BY, Ryan JJ. Hexabromocyclododecane (HBCD) in egg yolks from market bound Canadian chicken eggs, *Organohalogen Compounds* 2009;71: 1257 – 1261
- {RIVM}\_{ (The Netherlands Insitute for Fisheries Research). HBCD and TBBP-A in sewage sludge, sediments and biota, including interlaboratory study. Report No. C033/02. Authors: de Boer J, Allchin C, Zegers B, Boon JP, Brandsma SH, Morris S, Kruijt AW, van der Veen I, van Hesseligen JM, Haftka JHH. 2002
- ~~{RIVM}~~{RIVM (The Netherlands Insitute for Fisheries Research). Dietary intake of brominated flame retardants by the Duchth population. RIVM report 310305001/2003. Authors: de Winter-Sorkina R, Bakker MI, van Donkersgoed G, van Klaveren JD.
- Roos A, Nylund K, Häggberg L, Asplund L, Bergman A, Olsson M. Brominated flame retardants (BFR) in young grey seal males (*Halicoerus grypus*) from the Baltic Sea. Abstract from The Second International Workshop on Brominated Flame Retardants, May 14-16, Stockholm, Sweden. 2001; 337-341.
- Ronisz D, Finne EF, Karlsson H, Förlin L. Effects of the brominated flame retardants hexabromocyclododecane (HBCDD), and tetrabromobisphenol A (TBBPA), on hepatic enzymes and other biomarkers in juvenile rainbow trout and feral eelpout. *Aquat Toxicol.* 2004;69(3):229-45.
- Ryan, JJ, Wainman BC, Schecter A, Moisey J, Kosarac I and Sun WF . Trends of the brominated flame retardants, PBDEs and HBCD, in human milks from North America. *Organohalogen Compd.*, 2006;68, 778–781.
- Schlabach M, Fjeld E, Borgen AR. Brominated flame retardants in Drammens River and the Drammensfjord, Norway. Abstract, The Third International Workshop on Brominated Flame Retardants, June 6-9, Toronto, Canada. 2004a; 3779-3785.
- Schlabach M, Fjeld E, Gundersen H, Mariussen E, Kjellberg G, Breivik E. 2004b. Pollution of Lake Mjøsa by brominated flame retardants. *Organohalogen Compd* 2004b; 66: 3779-3785.
- Scottolini AG, Bhagavan NV, Oshiro TH, Abe SY. Serum high-density lipoprotein cholesterol concentrations in hypo- and hyperthyroidism. *Clin Chem.* 1980;26(5):584-7.
- Sellström U, Bignert A, Kierkegaard A, Häggberg L, de Wit CA, Olsson M, Jansson B. Temporal trend studies on tetra- and pentabrominated diphenyl ethers and hexabromocyclododecane in guillemot egg from the Baltic Sea. *Environ Sci Technol* 2003;37: 5496-5501.
- Seth R, Mackay D, Muncke J.. Estimating the organic carbon partition coefficient and its variability for hydrophobic chemicals. *Environ Sci Technol* 1999;33: 2390-2394.



- Smolarz K, Berger A. Long-term toxicity of hexabromocyclododecane (HBCDD) to the benthic clam *Macoma balthica* (L.) from the Baltic Sea. *Aquat Toxicol.* 2009;95 (3):239-47.
- Svobodova P M.Z. 2006. Brominated flame retardants in the environment: their sources and effects (a review). *Acta Vet. Brno* 2006; 75: 587-599.
- Swedish Environmental Protection Agency (Swedish EPA). Organic environmental pollutants in breast milk from Gothenburg, Sweden, 2001. Report 219 0108. Authors: Aune M, Barregård L, Claesson A, Darnerud PO. 2002, 10 pp.
- Swedish Environmental Protection Agency (Swedish EPA). Persistent organic pollutants (POP) in breast milk from primiparae women in Uppsala County, Sweden, 2002-2003. Livsmedelsverket, Uppsala, Sweden. Report 2150210 . Authors: Lignell S, Darnerud PO, Aune M, Törnkvist A. 2003, 9 pp.
- Swedish Environmental Protection Agency (Swedish EPA). Exponering för organiska miljökontaminanter via livsmedel. Livsmedelsverket, Uppsala, Sweden. Report 26. Authors: Lind Y, Darnerud PO, Aune M, Becker W. 2002; 103 pp.
- Sweeting, RM, Eales, GJ. Thyroxine 5'-monodeiodinase activity in microsomes from isolated hepatocytes of rainbow trout; effects of growth hormones and 3,5,3'-triiodo-L-thyronine. *Gen. Comp. Endocrinol.* 1992;88: 169-177.
- {TNO}. Hazardous chemicals in precipitation. TNO Environment, energy and Process Innovation. TNO Report R2003/198, 1-50. Netherlands, Netherlands Organization for Applied Scientific Research. 2003. Author: Peters RJB.
- {TNO}. Man-made chemicals in human blood. TNO Report R2004/493. <http://eu.greenpeace.org/downloads/chem/Blood-chemical-footprints.pdf> (2004). Author: Peters, RJB.
- Verreault J, Gabrielsen GW, Chu S, Muir DCG, Andersen M, Hamaed A, Letcher RJ. Flame retardants and methoxylated and hydroxylated polybrominated diphenyl ethers in two Norwegian Arctic top predators: glaucous gulls and polar bears. *Environ Sci Technol* 2005;39(16): 6021-6028.
- Verreault J, Gebbink WA, Gauthier LT, Gabrielsen GW, Letcher RJ. Brominated flame retardants in glaucous gulls from the Norwegian Arctic: more than just an issue of polybrominated diphenyl ethers. *Environ Sci Technol* 2007a.;41(14): 4925-4931.
- Verslycke TA, Vethaak AD, Arijs K, Janssen CR. Flame retardants, surfactants and organotins in sediment and mysid shrimp of the Scheldt estuary (The Netherlands). *Environ Pollut* 2005;136: 19-31.
- Viberg H, Fredriksson A, Eriksson P. Neonatal exposure to polybrominated diphenyl ether (PBDE 153) disrupts spontaneous behaviour, impairs learning and memory, and decreases hippocampal cholinergic receptors in adult mice. *Toxicol Appl Pharmacol.* 2003;192(2):95-106.
- Watanabe W, Shimizu T, Sawamura R, Hino A, Konno K, Hirose A, Kurokawa M. Effects of tetrabromobisphenol A, a brominated flame retardant, on the immune response to respiratory syncytial virus infection in mice. *Int Immunopharmacol.* 2010;10(4):393-7.
- {WIL}\_ (WIL Research Laboratories, Inc, Ashland, Ohio). A 28-day repeated dose oral toxicity study of HBCD in rats. USA. 1997; Author: Chengelis CP. 925 pp.
- {WSKOWWIN}\_ Water Solubility for Organic Compounds Program for Microsoft Windows {Estimation Model}\_ Version 1.41. Washington (DC): U.S. Environmental Protection Agency, Office of Pollution Prevention and Toxics; Syracuse (NY): Syracuse Research Corporation. 2000. Available from: [www.epa.gov/oppt/exposure/pubs/episuite.htm](http://www.epa.gov/oppt/exposure/pubs/episuite.htm)
- {WWF-UK}\_ (World Wildlife Fund, United Kingdom). National Biomonitoring Survey 2003, Thomson GO, Hodson S, Jones KC. Appendix 3: Lancaster University Analytica report, November 2003
- {WWF}\_ (World Wildlife Fund). Detox Campaign. Chemical check-up: An analysis of chemicals in the blood of members of European Parliament, April 2004;52-91. [http://www.wwf.dk/db/files/checkupmain\\_3.pdf](http://www.wwf.dk/db/files/checkupmain_3.pdf).
- Zennegg M, Kohler M, Gerecke AC, Schmid P. Polybrominated diphenyl ethers in whitefish from Swiss lakes and farmed rainbow trout. *Chemosphere* 2003;51: 545-553.
- Zennegg M, Brändli RC, Kupper T, Bucheli TD, Gujer E, Schmid P, Stadelmann FX, Tarradellas J. PCDD/Fs, PCBs, PBDEs, TBBPA and HBCD in compost and digestate. Abstract. *Dioxin* 2005, August 21-26, Toronto, Canada. p. 1040-1043.

Ziegler E, Anderson B, Haworth S, Lawlor T, Mortelmans K, Speck W. Salmonella mutagenicity tests III. Results from the testing of 255 chemicals. *Environ Mol Mutagen* 1987;9(suppl. 9):1-10.

Zitko V. Expanded polystyrene as a source of contaminants. *Marine Pollution Bulletin* 1993;26: 584-585.

---